

LYSIPILEX PLUS

DIETARY SUPPLEMENT

HELPS STIMULATE THE APPETITE**



WITH
VITAMIN C

SUGAR FREE

KramerNovis
Pharmaceuticals within reach of patients
www.kramernovis.com

Net Contents: 16 fl oz (473 mL)

Supplement Facts

Serving Size 1 tablespoonful (15 mL)

Servings Per Container 30

Amount Per Serving	%DV for adults and children 4 or more years of age
Vitamin A (as palmitate) 2750 mcg	306%
Vitamin C (as ascorbic acid) 45 mg	50%
Thiamine (as thiamine HCl) 50 mg	4167%
Riboflavin (vitamin B-2) 15 mg	1154%
Niacin (as niacinamide) 50 mg	313%
Vitamin B-6 (as pyridoxine HCl) 50 mg	2941%
Folate 1,333 mcg DFE (as 800 mcg folic acid)	333%
Vitamin B-12 (as cyanocobalamin) 50 mcg	2083%
Biotin 150 mcg	500%
Pantothenic Acid 50 mg (as calcium pantothenate)	1000%
Iron (as ferric ammonium citrate) 10 mg	56%
Zinc (as zinc sulfate) 15 mg	136%
Copper (as copper gluconate) 2 mg	222%
Molybdenum (as sodium molybdate) 15 mcg	33%
L-Lysine (as Lysine HCl) 1000 mg	*

*Daily Value not established.

Other ingredients: Purified water, glycerin, sorbitol, propylene glycol, PEG-40 hydrogenated castor oil, sucralose, citric acid, potassium sorbate (to preserve freshness), artificial and natural orange flavor, methylparaben (to preserve freshness) propylparaben (to preserve freshness), sodium bicarbonate.

Use: Dietary Supplement that helps stimulate the appetite**

Directions: Adults and children 4 years of age and older: Take one (1) tablespoonful (15 mL) preferably in the morning, as a dietary supplement, or as directed by your physician.

Warning: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

If you are under a physician's care or taking medication, contact your physician before using this product. **If pregnant or breast-feeding**, ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information: Store at controlled room temperature: 15°-30°C (59°-86°F). Avoid excessive heat.

Tamper Evident Feature: Do not use if inner seal is torn, broken or missing.

SHAKE WELL BEFORE USE

Manufactured in the USA for Kramer Novis, San Juan, PR 00917.

In case of an Adverse Event please call (787) 767-2072

www.kramernovis.com

**** This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.**

Item No. 52083-843-16

Lot No.
Exp. Date



Rev. 02/21