KEEP OUT OF REACH OF CHILDREN. Store in a cool, dry place.

For More Information: 1-888-462-2548

1-888-462-2548 TamaFlex.com

TamaFlex™ is a trademark of NXT USA, Inc. Patents Pending. AprèsFlex® is a registered

trademark of Laila Nutraceuticals exclusively licensed to PL Thomas -Laila Nutraceuticals LLC. U.S. Patent #8,551,496

Laila Nutraceuticals LLC. U.S. Patent #8,551,496 and other patents pending.



TAMAFLEX®

GNC

PROVEN JOINT SUPPORT

5 DAYS*

7x MORE COMFORT** & 7.5x BETTER FUNCTION** IN 30 DAYS

120 VEGETARIAN CAPSULES

60-DAY SUPPLY

ETARY SUPPLEMENT

TamaFlex*

CODE 304222

(Tamarind [Tamarindus indica] Seed Extract & Turmeric [Curcuma longa] Rhizome Extract)

AprèsFlex® *Boswellia serrata* Gum Extract

* Daily Value not established.

morning and one in the evening.

Serving Size One Capsule

Amount Per Serving

OTHER INGREDIENTS: Microcrystalline Cellulose, Vegetable Cellulose Capsule, Dicalcium Phosphate, Magnesium Stearate Vegetable Source, Calcium Silicate.

DIRECTIONS: As a dietary supplement, take one capsule in the

Supplement Facts

B)X(G

125 mg*

50 ma*

를 Distributed by: GNC Holdings, LLC, Pittsburgh, PA 15222 USA 望 Contains a bioengineered food ingredient.

통 WARNING: Consult your physician prior to using this product if you are pregnant, nursing, taking medication or have a medical condition. Discontinue use two weeks prior to surgery.

^ In a randomized, double-blinded trial, 90 healthy adults took 125 mg of TamaFlex™ or placebo twice daily for 90 days. Those taking TamaFlex™ had improved (i) joint comfort, stiffness and function (WOMAC), (ii) walking speed and distance and (ii) use of stairs.

(WOMAC), (ii) walking speed and distance and (iii) use of stairs. Improvements were seen as early as 2 weeks for comfort and stiffness. In a randomized, double-blind study, 67 adults took 50 mg of

In a randomized, double-blind study, of adults took 50 mg of AprèsFlex® or a placebo twice daily. Those taking AprèsFlex® had improvements in joint stiffness, comfort and function (WOMAC) that were 5, 7 and 43 times greater vs. the placebo group after 30 days. Improvements were seen as early as 5 days.