Store in a cool, dry place. For More Information

1-888-462-2548

SHOP NOW @ GNC.COM

Distributed by: General Nutrition Corporation Pittsburgh, PA 15222



GNC REVOND

RE-SHRED VASO-DRI COMPOUND

- + 5-DAY FORMULA DEGILLATES EXCESS WATER AS YOU STRIVE FOR A DRY,
- STRIATED PHYSIQUE* + FUELS INTENSE TRAINING TO DRIVE INSANE ROAD MAP VASCULARITY & MIND-BLOWING PUMPS WITH

VASODILATOR COMPOUND*

REPLENISH THE BODY PLUS INGREDIENTS FOR RAPID NUTRIENT **DELIVERY***

PHYSIOBOLIC ÆEDGE

DIETARY SUPPLEMENT **40 TABLETS** DIRECTIONS: As a dietary supplement, take four pills, twice daily. Consume 16oz

water with each dose Supplement Facts Serving Size Four Tablets Servings Per Container 10 Amount Per Serving % Daily Valu Magnesium (as Magnesium Oxide) . 25 ma.. Chloride (as Sodium Chloride 62 mg..

Sodium (as Sodium Chloride 25 mg.. Potassium (as Potassium Chloride) 25 mg. Vacodilator Compound L-Arginine I -Citrulline 500 ma Anmatine Sulfate 250 mg.... 300 mg.... 100 mg..... recVirta® trans. Resugratrol 30 mg.... **Illtra Water Complex** Dandelion Root Powder 1000 ma. Taraxacum officinalei

5	Amount Per Serving	% Daily Value	1
•	Parsley Leaf Powder	50 mg*	ı
	(Petroselinum crispum)		ı
	Horsetail Rush Stems Powder (Equisetum arvense)	50 mg	ı
ue	Hawthorn Leaves/Flowers Extrac	t 50 mg*	ı
%	(Crataegus laevigata)	F0 me *	ı
%	Cranberry Concentrate(Vaccinium macrocarpon)	50 mg	ı
%	Green Tea Leaf Extract	50 mg*	ı
	(Camellia sinensis) Elderberry Fruit Powder	20 ma *	ı
	(Sambucus nigra)		ı
	Buchu Leaves Extract	12.5 mg*	
	(Agathosma betulina) Sweet Clover Extract	10 ma *	ı
٠	(Malilatus afficinalis)		
	Corn Silk Powder (Zea mays)	10 mg*	
:	Uva Ursi Leaf Powder (Arctostaphylos uva ursi)	o mg	ı
	Juniper Berries Extract	* 5 mg*	ı
	(Juniperus communis)		1

Prickly Pear Fruit Extrac .50 mg......* (Oountia ficus-indical OTHER INGREDIENTS: Cellulose, Titanium Dioxide (Natural Mineral Whitener), Vegetable Acetoglycerides, Riboflavin (Color) WARNING: Consult your physician prior to using this product if you are pregnant, nursing,

Daily Value not established

aking medication, or have a medical condition. Discontinue use two weeks prior to surgery. These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.