

**TAMPER EVIDENT:** Do not use if imprinted seal under cap is missing or broken

**WARNING:** Accidental overdose of iron containing supplements is a leading cause of fatal poisoning in children under 6.

**KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.**

In case of accidental overdose, call a doctor or Poison Control Center immediately.

**WARNINGS:** Do not exceed recommended dosage. The treatment of any anemic condition should be under the supervision of a physician. Since oral iron products interfere with the absorption of oral tetracycline antibiotics, these products should not be taken within two hours of each other. This product may occasionally cause constipation, diarrhea or may color stool black. It may cause gastrointestinal discomfort (such as nausea) which may be minimized by taking with meals.

**As with any supplement,** If you are pregnant, nursing, or taking medication, ask a health professional before use.

Store at room temperature  
15° C - 30° C (59° - 86° F)



GERI  CARE®

**IRON**  
**27 MG**

**FERROUS GLUCONATE**

**IRON SUPPLEMENT**

**100 TABLETS**

## Supplement Facts

Serving Size 1 Tablet

	Amount Per Serving	%Daily Value
IRON (as Ferrous Gluconate)	27 mg	150%

Each tablet contains 27 mg of elemental iron equivalent to approximately 240 mg Ferrous Gluconate.

**Other ingredients:** Microcrystalline Cellulose, Croscarmellose Sodium, Stearic Acid, Hypromellose, Talc, Titanium Dioxide, Triacetin, Mineral Oil, FD&C Yellow #5 Lake (Tartrazine), Sodium Lauryl Sulfate, FD&C Blue #1 Lake.

**Suggested use:** Adults: take one (1) tablet daily with or after a meal as a dietary iron supplement, or as recommended by your physician.

Distributed By: **GERI-CARE PHARMACEUTICALS CORP.**

1295 Towbin Ave, Lakewood, NJ 08701

1-800-540-3765

REV 0723C