

Supplement Facts

Serving Size: 1 Capsule

	Amount Per Serving	%DV*
Ferrous Fumarate (anhydrous) (Equivalent to about 62.5 mg of Elemental Iron)	191.1 mg	
Polysaccharide Iron Complex (Equivalent to about 62.5 mg of Elemental Iron)	135.9 mg	** 694 %
Vitamin C (from ProAscorb C [†])	210 mg	350 %
Folic Acid	1 mg	250 %
Thiamine Mononitrate (B ₁)	5 mg	333 %
Riboflavin (B ₂)	5 mg	294 %
Niacin (B ₃)	20 mg	100 %
d-Calcium Pantothenate (B ₅)	7 mg	70 %
Pyridoxine HCl (B ₆)	25 mg	1250 %
Biotin (B ₇)	300 mcg	100 %
Cyanocobalamin (B ₁₂)	10 mcg	166 %

* Percent of Daily Value † U.S. Patent No: 5,626,883

** Total Iron: Ferrous Fumarate + Polysaccharide Iron Complex

Inactive Ingredients: Hypromellose, Silicon Dioxide, Magnesium Stearate, Titanium Dioxide, FDA/E172 Yellow Iron Oxide, FD&C Red #40, and FD&C Blue #1. Manufactured with Vcaps® capsule shells. Vcaps® and the Vcaps® Logo are trademarks used under license.

NDC 52747-712-60

Rx Only



IRON / FOLIC ACID / VITAMIN SUPPLEMENT CAPSULES

Store at controlled room temperature 15° to 30°C (59° to 86° F).
Keep in a cool, dry place. Capsules are not USP.



® Marketed by: US Pharmaceutical Corporation
Decatur, GA 30035, USA
Internet: www.uspco.com

90 Capsules

INDICATIONS: Integra Plus™ is indicated for the treatment of iron deficiency anemia and folate deficiency as in extended convalescence, menorrhagia, pregnancy, puberty, excessive blood loss and advanced age. Also for treatment of condition in which iron deficiency and vitamin C deficiency occur together, along with a deficient intake or increased need for B-Complex vitamins in chronic and acute illness, as well as cases of metabolic stress, and in convalescence.

DOSAGE AND ADMINISTRATION: Adults (persons over 12 years of age), one (1) capsule daily, between meals, or as prescribed by a physician. Do not exceed recommended dosage. Do not administer to children under the age of 12.

CAUTION: Rx only.

Rev. 01/2009

U.S. Patent No: 5,626,883

LC-10810

Consult package insert for full prescription information.

WARNING: Accidental overdose of iron-containing products is the leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or poison control center immediately.