



NDC 0904-5339-60

Prenatal Plus Iron Tablets

**Multivitamin/Multimineral
Supplement**For Use Before, During and
After Pregnancy.**Rx only****100 Tablets****Each Tablet Contains:
VITAMINS:**

	Amount per Tablet
Vitamin A (from Acetate and Beta Carotene)	4000 IU
Vitamin C (Ascorbic Acid)	120 mg
Vitamin D-3 (Cholecalciferol)	400 IU
Vitamin E (dl-Alpha Tocopheryl Acidate)	22 IU
Thiamine (Vitamin B-1 from Thiamine Mononitrate)	1.84 mg
Riboflavin (Vitamin B-2)	3 mg
Niacin (as Nicotinamide)	20 mg
Vitamin B-6 (from Pyridoxine HCl)	10 mg
Folic Acid	1 mg
Vitamin B-12 (Cyanocobalamin)	12 mcg

MINERALS:

Calcium (from Calcium Carbonate)	200 mg
Iron (from Carbonyl Iron)	29 mg
Zinc (from Zinc Oxide)	25 mg
Copper (from Cupric Oxide)	2 mg

Other Ingredients: Acetyl Polmalate, Citric Acid, hydroxytol, DL- alpha Tocopherol, Ethylcellulose, FD&C Blue #2 lake, FD&C Red #40 lake, FD&C Yellow #5 lake, FD&C Yellow #6 lake, Glucose, Gum Arabic, Hydroxypropyl Cellulose, Magnesium Stearate, Malt Starch, Maltodextrin, Methylcellulose, Microcrystalline Cellulose, Mineral Oil, Mono- and di-glycerides, Polyethylene Glycol, Pregelatinized Corn Starch, Silicon Dioxide, Sorbic Acid, Soy Potash, Stearic Acid, Sucrose, Talcum Oxide, Tricalcium Phosphate.

Indication: To provide Vitamin and Mineral supplementation throughout pregnancy and during the postnatal period for both the lactating and non-lactating mother. It is also useful for improving nutritional status prior to conception.

Dosage: As a dietary adjunct before, during and after pregnancy, take one tablet daily with a meal, or as directed by physician.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under age 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Precautions: Folic acid may partially correct the hematological damage due to Vitamin B12 deficiency of pernicious anemia, while the associated neurological damage progresses. This product contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

Store at room temperature. Dispense in a well closed light-resistant container, with a child resistant cap.

CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT SIDE EFFECTS. YOU MAY REPORT SIDE EFFECTS TO THE FDA AT 1-800-FDA-1088 (TOLL FREE)

Keep this and all medication out of the reach of children.

Do not use if imprinted Safety Seal under cap is broken or missing.

Manufactured by: Cepharma Inc, Cranbury, NJ 08512

Distributed by: MAJOR PHARMACEUTICALS, 31778 Enterprise Drive, Livonia, MI 48150 USA

Lot No. / Exp date Rev. 12/12

