

eNOXIDE™

Biomedical Nitric Oxide Technology


Fast, Potent NO Elevation

Sustains Maximum Pumps for Hours

Fastsorb™ Delivery

Patents Pending

The World's First Biomedical Super-Molecule



Delivers Real Nitric Oxide (NO)

Dietary Supplement
40 Fastsorb™ Tablets

WARNING: Never Exceed Recommended Dose.

MuscleMeds™

eNOXIDE

BIOMEDICAL SUPER-MOLECULE DELIVERS REAL NITRIC OXIDE

eNOXIDE is a revolutionary breakthrough in Nitric Oxide (NO) elevation made possible through the discovery and development of a biomedically engineered Real Nitric Oxide molecule administered in an advanced patent pending FASTSORB™ delivery tablet. eNOXIDE has the unique ability to deliver potent levels of actual NO molecules into the bloodstream and muscle tissue. This new Nitric Oxide Super-Molecule instantly triggers immense pumps, improves muscle performance and stimulates muscle growth. FASTSORB sub-buccal delivery of eNOXIDE's Real Nitric Oxide molecules ensure rapid, massively effective NO elevations within seconds and are sustained for hours.

eNOXIDE is a quantum leap above conventional NO products that merely attempt to increase NO levels using precursor ingredients such as Arginine AGK or Citrulline, which must go through the complex biochemical pathway to be converted to Nitric Oxide and these precursors only produce a small elevation in NO. eNOXIDE is not a nitric oxide precursor, its REAL NITRIC OXIDE!

eNOXIDE...Real Science... Real Nitric Oxide...Real Results!

- Keep out of reach of children.
- Store at 15°-30°C (59°-86°F).
- Protect from heat, light and moisture.
- Do not purchase if seal is broken.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Review warnings before using this product.

DIRECTIONS: eNOXIDE Fastsorb™ tablets are designed to be dissolved under your tongue. Ten minutes before your workout, place eNOXIDE Fastsorb™ tablet(s) under your tongue and let dissolve completely, then swallow. eNOXIDE should be dosed according to your body weight.

DOSAGE:

Body weight under 200 lbs. take 1 tablet.

Body weight over 200 lbs. take 1 to 2 tablet(s).

For healthy adults only. Do not exceed your recommended tablet dosage in a 12 hour period. Do not take more than 5 days per week. Maintain a level of good hydration while taking eNOXIDE.

Supplement Facts

Serving Size: 1 Tablet Servings Per Box: 40

	Amount Per Serving	% Daily Value
eNOXIN	10 mg	†
2-(nitrooxy)ethyl 2-amino-3-methylbutanoate		
† Daily value not established.		

Other ingredients: carboxymethylcellulose, lactose, phosphatidylcholine, phosphatidylserine, artificial flavor, aspartame, neotame, FD&C Red 40, FD&C Yellow 6, magnesium stearate and talc.

WARNING: NOT INTENDED FOR USE BY PERSONS WITH A MEDICAL CONDITION. NOT INTENDED FOR USE BY PERSONS UNDER AGE 18 OR PREGNANT OR LACTATING WOMEN. Consult a physician or licensed qualified healthcare professional before using this product. Not for people with cardiovascular diseases, blood pressure problems, head injury or other diseases or disorders. Consult your physician prior to use if you have a medical condition, including, but not limited to, heart, kidney, liver or thyroid disease, psychiatric or epileptic disorders, difficulty urinating, diabetes, high or low blood pressure, cardiac arrhythmia, recurrent headaches, enlarged prostate or glaucoma.

Do not take with any medication, especially cardiovascular and erectile dysfunction medications, blood pressure drugs, heart or other cardiovascular system medications, antibiotics, antihistamines, benzocaine and lidocaine or other drugs. Do not take with other supplement products designed to increase nitric oxide levels or products containing yohimbine or over 75 mg of niacin per serving. Avoid alcohol consumption within 5 hours before and after taking eNOXIDE.

Discontinue use immediately if you experience one or more of the following: dizziness, light headedness, severe headache, low blood pressure, rapid heartbeat, shortness of breath. Discontinue use 2 weeks prior to surgery. Do not use for more than 5 days per week, or more than 8 consecutive weeks.

USE ONLY AS DIRECTED. DO NOT EXCEED RECOMMENDED DAILY INTAKE.