
PEE



Powerful Weight Loss! A,1,2 Proven to Reduce BMI ▲,1,2

Fast-Acting Energizing Effects ♣,3 Clinically Proven Key Ingredients 1,2



Supplement Facts Serving Size: 2 Liquid-Caps Servings Per Container: 60	
Amount Per Serving % Daily V	alue
Vitamin C (as ascorbic acid) 7mg	12%
Folic acid 200mcg	50%
Calcium 50mg	5%
Iron (as ferrous gluconate dihydrate) 2mg	11%
Hydroxycut™ Max! Blend 456mg	†
Max! HydroxyBoost™	†
Caffeine anhydrous (1,3,7-trimethylxanthine)	t
L-alanine	Ť
L-serine	†
Max! ProDefine™	t
Alchemilla vulgaris extract (lady's mantle) (leaf)	<u>†</u> †
Olea europaea extract (wild olive) (leaf)	†
Cuminum cyminum extract (komijn) (seed)	<u> </u>
Mentha longifolia extract (wild mint) (leaf)	t
Max! Hydroxagen®	
Oleic acid	<u>†</u>
L-glutamic acid HCl	t
L-threonine	Ť
L-isoleucine	<u>†</u> †
Co-enzyme Q10	Ť
Glutathione	t
† Daily Value not established.	

OTHER INGREDIENTS: SESAME OIL CAPSULE (GELATIN, FROG RED NO. 40), TRICACIOUM PHOSPATE, CARAMEL OLOUR, INK, SPELIAC, TITANIUM DIODIE, AMMONIUM HYDROXIDE PROPULENE GLYCOL, POLYDIMETHY SI, OXANIE, GELATIN, POLYSOBRATE RO. CONTAINS SESAME, SULFITES AND WHEAT INGREDIENTS.

Product Information Line: 1-877-502-1998 ◆◆Based on IRI F/D/MX sales data for Hydroxycut™ caplets 2007-2008.

DIRECTIONS: To assess individual tolerance, for the first two days, start with one (1) Liquid-Cap taken with a glass of water two (2) times daily, approximately 30 minutes before your main meals. For days three and beyond, follow the guidelines below 0 not exceed 2 Liquid-Caps in a 4-hour period and/or 6 Liquid-Caps in a 24-hour period. Always meal the entire label before use, and follow the directions provided.

New to Pro Clinical Hydroxycut™ Max?
To assess individual tolerance, follow these guidelines:

• Day 1 and 2: Take 1 Liquid-Cap, 2x daily.

• Day 3 and 4: Take 2 Liquid-Caps, 2x daily.

• Day 5 and Beyond: Take 2 Liquid-Caps, 3x daily.

For Best Results:

• Use for 8-12 weeks with diet and exercise.

• Stay hydrated by drinking 8-10 glasses of water per day. Do not snack between meals.
 Do not take within 5 hours of bedtime.

WARNING: Not intended for use by persons under 18. Do not use if pregnant or rursing. Accidental overdose of iron-containing products is a leading cause of ratal poisoning in children under 6. in case of accident overdose, call a medical doctor or poison control immediately. Take a few hours before or after taking other medications. Discontinue use and consult a medical doctor if you experience unusual symptoms. Consult a medical doctor before use if you have been treated for, diagnosed with or have a doctor legicle use in your rare usern analist on, diagnoss must be family history of any medical condition of you are using any prescription or over-the-counter drugs), including blood thinners. One serving (2 capsules) of this product contains up to as much cafferine as 3 cups of coffee (270mg). Cafferine sensitive individuals may experience the following symptoms: camenie sensitive inioviduolis indegreparente die following symptoms inioluding (but not limited to) estellessness, nervousness, tremors, headache, anxiety, palpitations, increased heart rate or difficulty sleeping. Do not combine with other sources of caffeine. Consult a medical doctor before starting any diet or exercise plan. Do not exceed recommended serving stating any une or belocks plan to be forecast execution menues serving. Improper use of this product will not improve results and is not advised. Use only as directed. **Do not use if packaging has been tampered with.** Store in a cool, dry place (60°F to 80°F). **KEEP OUT OF REACH OF CHILDREN.**

*These statements have not been evaluated by the Food and Drug Administration This product is not intended to diagnose, treat, cure, or prevent any disease.

Made in the U.S.A. from international and domestic ingredients. Distributed by lovate Health Sciences USA, Inc., 1106 North Market Street, Suite 1330, Wilmington, DE 19801. © 2010. Capsule shown not actual size. For lot no, and expiry date: see bottle

New Pro Clinical Hydroxycut™ Max! - It Really Works! New Pro Clinical Hydroxycut™ Max! delivers more of what women want in

a weight-loss solution.

Weight-Loss Ingredients
Proven in Two Clinical Studies 1,2 Fast-Acting Energizing Effects 4,3 Powerful Weight Loss 4,1,2

Female-Friendly Ingredients: · Folic Acid & Iron

Two Clinical Studies: Subjects Lost 8x the Weight and Reduced BMI (Body Mass Index).

Data analysis of the results from both clinical trials on the key ingredients in Pro Clinical Hydroxycut™ Max! shows there is statistical significance for test subjects losing 7 times more weight versus placebo subjects.

1 Study one: In a double-blind, placebo-controlled, clinical trial conducted by third-party researchers, individuals taking the key ingredient combination (Max! ProDefine™, Alchemilla vulgaris, Olea europaea, Cuminum cyminum, Mentha longifolia) lost an average of 20.94 lbs. over 12 weeks, compared to those using a placebo who lost an average of 1.70 lbs. Test subjects also significantly reduced BMI versus the placebo group (10.2% vs 0.9%). Both groups followed a calorie-reduced diet.

2 Study two: In a double-blind, placebo-controlled, clinical trial funded by the makers of Hydroxycut™ Max! and conducted by third-party researchers, individuals taking the same key ingredient combination lost an average of 16.50 lbs. in 8 weeks, compared to those taking a placebo, who lost an average of only 1.73 lbs. Test subjects also significantly reduced BMI versus the placebo group (8.1% vs 0.8%). Both groups followed a calorie-reduced diet. A

Fast-Acting Energizing Effects[♠]

3 New Pro Clinical Hydroxycut™ Max! contains an ingredient (caffeine anhydrous [1,3,7-trimethylxanthine]) that supports increased energy.▲