



CAVALIER™

GENTLEMEN'S PRODUCT

**DAILY
PROSTATE
SUPPORT®**

FOR GENTLEMEN OF
HIGH DISTINCTION

90
CAPSULES

EXCLUSIVELY
TAILORED
FORMULA

DIETARY SUPPLEMENT

SUGGESTED USE:
TAKE ONE SERVING (3 CAPSULES)
ONCE DAILY WITH FOOD.
DO NOT EXCEED 6 CAPSULES IN A 24-HOUR
PERIOD. USE FOR UP TO 8 WEEKS IN A CYCLE.
THEN DISCONTINUE USE FOR 4 WEEKS
BEFORE STARTING A NEW CYCLE.

SUPPLEMENT FACTS

Serving Size: 3 Capsules

Servings Per Container: 30

	AMOUNT PER SERVING	%DV**
Vitamin D (as Cholecalciferol)	841 IU	210%
Zinc (as Zinc Monomethionine)	30 mg	200%

Cavalier™ Prostate Blend	1423 mg	†
Pomegranate (<i>Punica granatum</i>) fruit extract, Saw Palmetto (<i>Serenoa repens</i>) fruit extract, Lycopene 10% Beadlets, Pumpkin (<i>Cucurbita moschata</i>) seed extract, Japanese sophora (<i>Sophora japonica</i>) bud extract (standardized for Quercetin), Soy bean (<i>Glycine max</i>) seed extract (standardized for Beta Sitosterol), Pygeum (<i>Prunus africana</i>) bark		

**Percent Daily Values are based on a 2,000 calorie diet.

† Daily Value not established.

OTHER INGREDIENTS:

Microcrystalline Cellulose, Silicon Dioxide, Magnesium Stearate, Gelatin, Titanium Dioxide, FD&C Blue #1, FD&C Red #40.

CONTAINS: SOY

THESE STATEMENTS HAVE NOT BEEN
EVALUATED BY THE FOOD AND DRUG
ADMINISTRATION. THIS PRODUCT IS NOT
INTENDED TO DIAGNOSE, TREAT, CURE,
OR PREVENT ANY DISEASE.

 **nutrabolt™**

DISTRIBUTED BY:
NUTRABOLT™

3891 S. TRADITIONS DR.
BRYAN, TX 77807

100994



WARNING: THIS PRODUCT IS ONLY INTENDED FOR HEALTHY ADULT MEN, 18 YEARS OF AGE OR OLDER. NOT FOR USE BY WOMEN, ESPECIALLY THOSE WHO ARE PREGNANT OR NURSING. CONSULT A LICENSED, QUALIFIED HEALTH CARE PROFESSIONAL BEFORE USING THIS PRODUCT, INCLUDING BUT NOT LIMITED TO, IF YOU HAVE BEEN TREATED FOR, DIAGNOSED WITH, OR HAVE A FAMILY HISTORY OF ANY MEDICAL CONDITION, INCLUDING BUT NOT LIMITED TO: PROSTATE CANCER OR PROSTATE ENLARGEMENT, OR, IF YOU ARE TAKING ANY OTHER DIETARY SUPPLEMENT, PRESCRIPTION DRUG OR OVER-THE-COUNTER MEDICATION. DO NOT USE THIS PRODUCT IF YOU HAVE ANY SERIOUS MEDICAL CONDITIONS. DISCONTINUE USE 2 WEEKS PRIOR TO SURGERY. IMMEDIATELY DISCONTINUE USE AND CONSULT A LICENSED, QUALIFIED HEALTH CARE PROFESSIONAL IF YOU EXPERIENCE ANY ADVERSE REACTION TO THIS PRODUCT. DO NOT EXCEED RECOMMENDATIONS FOR SUGGESTED USE. DO NOT USE IF SAFETY SEAL IS BROKEN OR MISSING. STORE IN A COOL, DRY PLACE. KEEP OUT OF REACH OF CHILDREN.