d water, C Blue #1.	Amount per Serving: Serving Size: 1 Softgel	
	Vitamin C (as ascorbic acid)	30 mg
FD&C	Biotin	75 mc
a di a	Vitamin D3 (as cholecalciferol)	500 IL
	Vitamin B12 (as cyanocobalamin)	13 mcg
stramel and so	Folate (as ((6S)-N5-methyl-	1 mg
5 5 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	tetrahydrofolic acid calcium salt	
를 <b>불</b> 물 돌	(molar equivalent to 600 mcg of folic acid	)
13 E 20	and folic acid, USP 400 mcg)	8
BC F	Iron (as ferrous asparto glycinate) Sumalate*	10 mg
FD8(	lodine (as potassium iodide)	150 mcg
돌를	Vitamin B6 (as pyridoxine HCl)	5 mg
8.8	Vitamin E (as di-alpha tocopheryl acetate)	10
8 E E	Docosahexaenoic acid (DHA)	200 mg
88 E E	Blueberry extract (vaccinium spp.)	5 mg



DOSAGE AND ADMINISTRATION: One softgel daily, or as directed by a physician. See package outsert for full prescribing information.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

## KEEP OUT OF REACH OF CHILDREN.

STORAGE: Store at 20°-25°C (68°-77°F); excursions permitted to 15° - 30°C (59°-86°F) ISee USP Controlled Room Temperature.1

Lot# Exp:

rev. 0814