may very well be the master key to

Think about it this way ... If Testosterone is the pedal to the gas, then Myostatin is the brake. Myostatin decreases muscle growth and differentiation - regulate Myostatin downward, and you may very well be on the way to increasing your

Glyco-proteins are uniquely structured compounds, which are responsible for an array of activites, including hormone regulation. KING™ is based on

promising research surrounding glyco-proteins and the role of myostatin

in muscle-building. This allows us the ability to roll this scientific data into one

"SUPER PILL" KING™



Manufactured for and Distributed by: CUTLER NUTRITION Hollywood, FL 33312 To report an adverse event or for more information call: 954,926,0900 (tel) www.cutlernutrition.com

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE. TREAT.

Statements based on early-stage independent 3rd party in vivo and / or in vitro model scientific research data findings for individual



mtor anabolic signaling agent*†

CLINICALLY RESEARCHED INGREDIENTS

PROMOTES IGF-2 EXPRESSION

BOOSTS MYOGENIC REGULATORY FACTORS (MRFS)*

SUPPORTS MUSCLE FIBERS®

MAY PROMOTE:

ENHANCED STRENGTH** LEAN MUSCLE MASS*1 ANABOLIC ACTIVITY**



DIETARY SUPPLEMENT

Supplement Facts

Serving Size 2 capsules Servings Per Container 30

Amount Per Serving

% Daily Value

Vitamin D3 (as cholecalciferol)

4.800 IU 1.200% DV

GLYCOPROTEIN MUSCLE GROWTH BLEND (proprietary)

Activator™ (Phosphatidic Acid {Lysophosphatidic Acid, Phosphatidylserine, Phosphatidylcholine, Phosphatidylethanolamine. Phosphotidylinositiol3)

Niagen™ (as Nicotinamide riboside)

** Daily Value (DV) not established

Other Ingredients: Gelatin, microcrystalline cellulose. maltodextrin, silica, cellulose (as ethyl cellulose), magnesium stearate, stearic acid, FD&C Blue No. 1, FD&C Red No. 40. FD&C Yellow No. 6 and titanium dioxide.

Contains sov.

Niagen™ is a registered trademark of Chromadex, inc.

Please read entire label before use.


MAGESTED USE:

Take two (2) capsules daily with food, or as recommended by a healthcare practitioner.



Not intended for use by persons under age 18. Do not exceed recommended dose. Do not take for more than eight (8) consecutive weeks. Get the consent of a licensed physician before using this product, especially if you are taking medication or have a medical condition. This product should not be taken by women. Do not take if you are pregnant, lactating or trying to become pregnant. Do not use if allergic or contraindicated to aspirin. Discontinue use two weeks prior to surgery or if stomach upset occurs. KEEP THIS PRODUCT AND ALL SUPPLEMENTS OUT OF THE REACH OF CHILDREN.





During development, mesodermal stem cells become committed to myogenic fate. Muscle precursors (myoblasts) remain in a proliferative state until they exit from the cell cycle and are differentiate. instructed Differentiation is accompanied by fusion, where committed myoblasts become polynucleated myotubes.

