



**HI-TECH
PHARMACEUTICALS**
Dietary Supplement

FASTIN®

(Phenylethylamine HCl)

• **Pharmaceutical-Grade
Weight Loss Aid for
Optimal Diet & Energy!†**

60 Tablets

345mg

Supplement Facts

Serving Size: 1 Tablet **Servings per Container:** 60

| | Amount Per Serving | % Daily Value |
|---|--------------------|---------------|
| Proprietary blend with Thermo-Rx®: * Phenylethylamine HCl, Methylsinephrine HCl, Theobromine Anhydrous, 1,3-Dimethylamine HCl, Synephrine HCl, N-Methyl-B-Phenylethylamine HCl, Yohimbine HCl | 245mg | * |
| Caffeine (anhydrous) | 100mg | * |

*Daily value not established

Other Ingredients: Dextrose, Microcrystalline Cellulose, Hydroxypropyl Methylcellulose, Stearic Acid, Magnesium Stearate, Sodium Starch Glycolate, Starch, Silica.

Directions: Take 1-2 tablets in the morning and 1 tablet after lunch. Do not exceed 4 tablets daily. Fastin® is heat and moisture sensitive, and the bottle should remain sealed after using. Keep desiccant in the bottle to avoid moisture. **Store at room temperature 59° - 86°F protected from moisture, heat, and light.** Failure to do so may cause pills to slightly turn brown. This will not affect the efficacy, potency or safety of the product, but will cause Fastin® to become brittle.

Manufactured by: Hi-Tech Pharmaceuticals, Inc.

6015-B Unity Drive • Norcross GA 30071 • 1.888.855.7919

†THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT ANY DISEASE.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF YOU ARE PREGNANT OR NURSING. THIS PRODUCT CAN RAISE BLOOD PRESSURE AND INTERFERE WITH OTHER DRUGS YOU MAY BE TAKING. TALK TO YOUR DOCTOR ABOUT THIS PRODUCT.

(WARNINGS/INFORMATION CONTINUED ON BACK OF LABEL) SALE TO PERSONS 17 YEARS OF AGE OR YOUNGER IS PROHIBITED.

INDIVIDUALS WHO CONSUME CAFFEINE WITH THIS PRODUCT MAY EXPERIENCE SERIOUS ADVERSE HEALTH EFFECTS. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A LICENSED HEALTH CARE PROFESSIONAL BEFORE CONSUMING THIS PRODUCT. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A DOCTOR BEFORE CONSUMING THIS PRODUCT. DO NOT CONSUME SYNEPHRINE OR CAFFEINE FROM OTHER SOURCES, INCLUDING BUT NOT LIMITED TO, COFFEE, TEA, SODA OR OTHER DIETARY PRODUCTS OR MEDICATIONS CONTAINING CAFFEINE OR PHENYLEPHRINE. DO NOT USE FOR MORE THAN 8 WEEKS. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU ARE TAKING MEDICATION, INCLUDING BUT NOT LIMITED TO MAOI INHIBITORS, ASPIRIN, ANTIDEPRESSANTS, NONSTEROIDAL ANTI-INFLAMMATORY DRUGS OR PRODUCTS CONTAINING PHYLEPHRINE, EPHEDRINE, PSEUDOEPHEDRINE, OR OTHER STIMULANTS. CONSULT YOUR PHYSICIAN PRIOR TO USE IF YOU HAVE A MEDICAL CONDITION, INCLUDING BUT NOT LIMITED TO, HEART, LIVER, KIDNEY, OR THYROID DISEASE, PSYCHIATRIC OR EPILEPTIC DISORDERS, DIFFICULTY URINATING, DIABETES, HIGH BLOOD PRESSURE, CARDIAC ARRHYTHMIA, RECURRENT HEADACHES, ENLARGED PROSTATE OR GLAUCOMA. DISCONTINUE 2 WEEKS PRIOR TO SURGERY OR IF YOU EXPERIENCE RAPID HEART BEAT, DIZZINESS, SEVERE HEADACHE OR SHORTNESS OF BREATH. IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY. AVOID ALCOHOL WHILE TAKING THIS PRODUCT. EXCEEDING RECOMMENDED SERVING MAY CAUSE SERIOUS ADVERSE HEALTH EFFECTS, INCLUDING HEART ATTACK AND STROKE. IMPROPER USE OF THIS PRODUCT MAY BE HAZARDOUS TO A PERSON'S HEALTH.

**TO REPORT ADVERSE EFFECTS CALL FDA'S
MEDWATCH AT 1-800-332-1088**



8 57084 00074 3

Tamper Evident
Do Not Use If Printed Seal Is
Cut, Torn Or Missing

**PEEL
HERE**