



Dietary Supplement

Stimerex[®] HARDCORE

The Ultimate Diet & Energy Aid
Developed for Hardcore Athletes!†

The World's Strongest Stimulant with
Acacia Rigidula and 1,3-Dimethylamylamine†

SOCT

SUPPLEMENT FACTS

Serving Size: 1 capsule Servings Per Container: 90

Amount Per Serving % Daily Value

Proprietary Blend: * 475mg
Acacia Rigidula 98% Extract (leaves), [Yielding R-Beta-Methylphenylethylamine, N-Methyl-B-Phenylethylamine, B-Phenylethylamine, and Methylsynephrine], Theobromine, 1,3-Dimethylamylamine HCl, Green Tea Extract (leaves), Naringen (fruit), 3-Dimethyl Aminoethanol Bitartrate, Yohimbe Extract (bark)

Caffeine (anhydrous) * 150mg

*Daily value not established

Other Ingredients: Dicalcium Phosphate, Sodium Starch Glycolate, Magnesium Stearate, Silica, FD&C Blue #1, FD&C Red #40, FD&C Yellow #6

Dosage: Take 1 capsule three times daily. Do not exceed 4 capsules in any 24-hour period.



Manufactured by: Hi-Tech Pharmaceuticals, Inc.
6015-B Unity Drive • Norcross • GA • 30071
www.hitechpharma.com • 1.888.855.7919

†THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT ANY DISEASE.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF YOU ARE PREGNANT OR NURSING. THIS PRODUCT CAN RAISE BLOOD PRESSURE AND INTERFERE WITH OTHER DRUGS YOU MAY BE TAKING. TALK TO YOUR DOCTOR ABOUT THIS PRODUCT.

(WARNINGS/INFORMATION CONTINUED ON BACK OF LABEL)
SALE TO PERSONS 17 YEARS OF AGE OR YOUNGER IS PROHIBITED.



Tamper Evident
Do Not Use if Printed
Seal Is Cut, Torn
Or Missing

PEEL
HERE

**TO REPORT ADVERSE EFFECTS CALL FDA'S
MEDWATCH AT 1-800-332-1088**

INDIVIDUALS WHO CONSUME CAFFEINE WITH THIS PRODUCT MAY EXPERIENCE SERIOUS ADVERSE HEALTH EFFECTS. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A LICENSED HEALTH CARE PROFESSIONAL BEFORE CONSUMING THIS PRODUCT. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A DOCTOR BEFORE CONSUMING THIS PRODUCT. DO NOT CONSUME SYNEPHRINE OR CAFFEINE FROM OTHER SOURCES, INCLUDING BUT NOT LIMITED TO, COFFEE, TEA, SODA OR OTHER DIETARY PRODUCTS OR MEDICATIONS CONTAINING CAFFEINE OR PHENYLEPHRINE. DO NOT USE FOR MORE THAN 8 WEEKS. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU ARE TAKING MEDICATION, INCLUDING BUT NOT LIMITED TO MAOI INHIBITORS, ASPIRIN, ANTIDEPRESSANTS, NONSTEROIDAL ANTI-INFLAMMATORY DRUGS OR PRODUCTS CONTAINING PHENYLEPHRINE, EPHEDRINE, PSEUDOEPHEDRINE, OR OTHER STIMULANTS. CONSULT YOUR PHYSICIAN PRIOR TO USE IF YOU HAVE A MEDICAL CONDITION, INCLUDING BUT NOT LIMITED TO, HEART, LIVER, KIDNEY, OR THYROID DISEASE, PSYCHIATRIC OR EPILEPTIC DISORDERS, DIFFICULTY URINATING, DIABETES, HIGH BLOOD PRESSURE, CARDIAC ARRHYTHMIA, RECURRENT HEADACHES, ENLARGED PROSTATE OR GLAUCOMA. DISCONTINUE 2 WEEKS PRIOR TO SURGERY OR IF YOU EXPERIENCE RAPID HEART BEAT, DIZZINESS, SEVERE HEADACHE OR SHORTNESS OF BREATH. IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY. AVOID ALCOHOL WHILE TAKING THIS PRODUCT. EXCEEDING RECOMMENDED SERVING MAY CAUSE SERIOUS ADVERSE HEALTH EFFECTS, INCLUDING HEART ATTACK AND STROKE. IMPROPER USE OF THIS PRODUCT MAY BE HAZARDOUS TO A PERSON'S HEALTH.