



Lipodrene[®] XTREME

DIETARY SUPPLEMENT 645MG

EXTREME WEIGHT LOSS
& ENERGY AID†

REVOLUTIONARY DIET & ENERGY AID WITH
ISOPROPYLNORSYNEPHRINE AND CITRUS-RX!†

90CT

Supplement Facts

Serving Size: 1 Tablet Servings Per Container: 90

Amount Per Serving %Daily Value

Proprietary Blend: 395mg *

Isopropylorsynephrine HCl, Acacia Rigidula Extract 95% (leaves)(yielding 175mg Phenylethylamine Alkaloids), Caffeine Anhydrous, 1,3-Dimethylamylamine HCl, Green Tea 98% Extract (leaves)(standardized to EGCG, Theophylline, and Polyphenols), Theobromine Anhydrous, Yohimbe Extract (bark), Hoodia Gordonii Extract (cactus), Naringen (fruit), 5-Methoxytryptamine HCl, L-5-Hydroxytryptophan, Raspberry Ketones

Citrus-Rx™ Proprietary Lipolytic Blend: 250mg *

Citrus Natsudaikai Hayata Extract (fruit), Citrus Junos Sieb. Ex Tanaka Extract (fruit), Citrus limonium Extract(fruit), Citrus Aurantium Extract (fruit)(supplying 25mg Synephrine Alkaloids)

*Daily value not established

Other Ingredients: Dextrose, Microcrystalline Cellulose, Hydroxypropyl Methylcellulose, Dicalcium Phosphate, Sodium Starch Glycolate, Magnesium Stearate, Stearic Acid, Povidone, Silica, FD&C Red #40

Suggested Dose: Take 1-2 tablets in the morning and 1 tablet in the afternoon. Do not exceed recommended dose.



Hi-Tech Pharmaceuticals, Inc.
6015-B Unity Dr • Norcross, GA 30071
www.hitechpharma.com • 1.888.855.7919

†THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT ANY DISEASE.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF YOU ARE PREGNANT OR NURSING. THIS PRODUCT CAN RAISE BLOOD PRESSURE AND INTERFERE WITH OTHER DRUGS YOU MAY BE TAKING. TALK TO YOUR DOCTOR ABOUT THIS PRODUCT.

**(WARNINGS/INFORMATION CONTINUED ON BACK OF LABEL)
SALE TO PERSONS 17 YEARS OF AGE OR YOUNGER IS PROHIBITED.**

PEEL
HERE
Tamper Evident
Do Not Use If Printed Seal is
Cut, Torn Or Missing



INDIVIDUALS WHO CONSUME CAFFEINE WITH THIS PRODUCT MAY EXPERIENCE SERIOUS ADVERSE HEALTH EFFECTS. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A LICENSED HEALTH CARE PROFESSIONAL BEFORE CONSUMING THIS PRODUCT. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A DOCTOR BEFORE CONSUMING THIS PRODUCT. DO NOT CONSUME SYNEPHRINE OR CAFFEINE FROM OTHER SOURCES, INCLUDING BUT NOT LIMITED TO, COFFEE, TEA, SODA OR OTHER DIETARY PRODUCTS OR MEDICATIONS CONTAINING CAFFEINE OR PHENYLEPHRINE. DO NOT USE FOR MORE THAN 8 WEEKS. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU ARE TAKING MEDICATION, INCLUDING BUT NOT LIMITED TO MAOI INHIBITORS, ASPIRIN, ANTIDEPRESSANTS, NONSTEROIDAL ANTI-INFLAMMATORY DRUGS OR PRODUCTS CONTAINING PHENYLEPHRINE, EPHEDRINE, PSEUDOEPHEDRINE, OR OTHER STIMULANTS. CONSULT YOUR PHYSICIAN PRIOR TO USE IF YOU HAVE A MEDICAL CONDITION, INCLUDING BUT NOT LIMITED TO, HEART, LIVER, KIDNEY, OR THYROID DISEASE, PSYCHIATRIC OR EPILEPTIC DISORDERS, DIFFICULTY URINATING, DIABETES, HIGH BLOOD PRESSURE, CARDIAC ARRHYTHMIA, RECURRENT HEADACHES, ENLARGED PROSTATE OR GLAUCOMA. DISCONTINUE 2 WEEKS PRIOR TO SURGERY OR IF YOU EXPERIENCE RAPID HEART BEAT, DIZZINESS, SEVERE HEADACHE OR SHORTNESS OF BREATH. IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY. AVOID ALCOHOL WHILE TAKING THIS PRODUCT. EXCEEDING RECOMMENDED SERVING MAY CAUSE SERIOUS ADVERSE HEALTH EFFECTS, INCLUDING HEART ATTACK AND STROKE. IMPROPER USE OF THIS PRODUCT MAY BE HAZARDOUS TO A PERSON'S HEALTH.

**TO REPORT ADVERSE EFFECTS CALL FDA'S
MEDWATCH AT 1-800-332-1088**