

HI-TECH PHARMACEUTICALS, INC.

Dietary Supplement

FASTIN[®]
(Phenylethylamine HCL)

The World's Most Advanced
Weight Loss Aid Ever Developed!†

EXP: 05/17



HI-TECH PHARMACEUTICALS, INC.



RAPID FAT BURNER†

Dietary Supplement

FASTIN[®]

- Increases the metabolic rate, promoting thermogenesis (The Burning of Stored Body Fat)†

- Increases the release of Norepinephrine and Dopamine for dramatic weight loss†

- Promotes extreme energy and does not cause the "jitters"†



References:

- (1) Beneficial Effects of Fastin[®] on Weight Loss, Energy Expenditure, and Fat Loss; Pharmatech Ltd. BZ, C.A. 2006
- (2) Am J Med. 2005 Sept; 118(9):998-1003
- (3) Metabolism 1991 40(3):323-329
- (4) J Med. 2002; 33(1-4):247-264
- (5) Med Hypotheses 2002 June; 58(6):491-5

RAPID FAT LOSS CATALYST†

THERMOGENIC INTENSIFIER†

RAPID FAT BURNER†

Dietary Supplement

FASTIN[®]

(Phenylethylamine HCL)



RAPID FAT LOSS
THERMOGENIC INTENSIFIER†



30 CT

THERMOGENIC INTENSIFIER†

RAPID FAT BURNER†

- Increases the metabolic rate, promoting thermogenesis (The Burning of Stored Body Fat)†

- Increases the release of Norepinephrine and Dopamine for dramatic weight loss†

- Promotes extreme energy and does not cause the "jitters"†



The World's
Most Advanced
Weight Loss Aid
Ever Developed!†

THERMOGENIC INTENSIFIER†

RAPID FAT BURNER†

Supplement Facts

Serving Size: 1 Tablet

Servings Per Container: 30

Amount Per Serving % Daily Value

Proprietary blend with Thermo-Rx[®]: 245mg
Phenylethylamine HCl, Methylsynephrine HCl,
Theobromine Anhydrous, 1,3-Dimethylamine HCl,
Synephrine HCl, N-Methyl-B-Phenylethylamine HCl,
Yohimbine HCl

Caffeine (Anhydrous) 100mg

*Daily value not established

Other Ingredients: Dextrose, Microcrystalline Cellulose, Hydroxypropyl Methylcellulose, Stearic Acid, Magnesium Stearate, Sodium Starch Glycolate, Starch, Silica.

Dosage: Take 1-2 tablets in the morning and 1 tablet after lunch. Do not exceed 4 tablets daily. Fastin[®] is heat and moisture sensitive, and the bottle should remain sealed after using. Keep desiccant in the bottle to avoid moisture. **Store at room temperature 59°-86°F protected from moisture, heat, and light.** Failure to do so may cause pills to turn slightly brown. This will not affect the efficacy, potency or safety of the product, but will cause Fastin[®] to become brittle.

Manufactured & Distributed By: Hi-Tech Pharmaceuticals, Inc. 6015-B Unity Drive • Norcross, GA 30071 • 1.888.855.7919

†THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT ANY DISEASE.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF YOU ARE PREGNANT OR NURSING. INDIVIDUALS WHO CONSUME CAFFEINE WITH THIS PRODUCT MAY EXPERIENCE SERIOUS ADVERSE HEALTH EFFECTS. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A LICENSED HEALTH CARE PROFESSIONAL BEFORE CONSUMING THIS PRODUCT. IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY. AVOID ALCOHOL WHILE TAKING THIS PRODUCT. DO NOT EXCEED RECOMMENDED SERVING. EXCEEDING RECOMMENDED SERVING MAY CAUSE SERIOUS ADVERSE HEALTH EFFECTS, INCLUDING HEART ATTACK AND STROKE. DISCONTINUE USE AND CALL A PHYSICIAN OR LICENSED QUALIFIED HEALTH CARE PROFESSIONAL IMMEDIATELY IF YOU EXPERIENCE RAPID HEARTBEAT, DIZZINESS, SEVERE HEADACHE, SHORTNESS OF BREATH, OR OTHER SIMILAR SYMPTOMS. IMPROPER USE OF THIS PRODUCT MAY BE HAZARDOUS TO A PERSON'S HEALTH. EXCEEDING RECOMMENDED SERVING WILL NOT IMPROVE RESULTS. TO REPORT ADVERSE EFFECTS CALL FDA'S MEDWATCH AT 1.800.332.1088. DO NOT USE FOR MORE THAN 8 WEEKS. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU HAVE A MEDICAL CONDITION, INCLUDING BUT NOT LIMITED TO, HEART, LIVER, KIDNEY OR THYROID DISEASE, PSYCHIATRIC OR EPILEPTIC DISORDERS, DIFFICULTY URINATING, DIABETES, HIGH BLOOD PRESSURE, CARDIAC ARRHYTHMIA, RECURRENT HEADACHES, ENLARGED PROSTATE OR GLAUCOMA. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU ARE TAKING MEDICATION, INCLUDING BUT NOT LIMITED TO, MAOI INHIBITORS, ANTIDEPRESSANTS, ASPIRIN, NONSTEROIDAL ANTI-INFLAMMATORY DRUGS OR PRODUCTS CONTAINING PHENYLEPHRINE, EPHEDRINE, PSEUDOEPHEDRINE, OR OTHER STIMULANTS. DISCONTINUE USE TWO WEEKS PRIOR TO SURGERY.

THERMOGENIC INTENSIFIER†