## LIPORUSH® DS

## Supplement Facts

Serving Size 1 Capsule Servings Per Container 60 Servings

Amount Per Serving % DV Lipo|Burn Blend: 435 mg Caffeine Anhydrous (300 mg). Amla (Phyllanthus emblica)(Fruit) Extract Higenamine HCl, Hordenine.

Synephrine HCI Dendrobium nobile (Stem) Concentrate, Theobromine.

Yohimbine HCI (17a-Hydroxy-Yohimban-16a-Carboxylic Acid Methyl Ester), Alpha Yohimbine (Rauwolscine)

Water|Shred Blend: 175 mg Taraxacum officianales (Dandelion Root) Extract 4:1 (1009) Arctostaphylos pincuica (Uva Ursi Leaf Powder) Commiphora mukul Gugulipid® Extract (Standardized to 2.5% Guggulsterones), L-Carnitine

Neuro Recovery Blend: 50 ma Chinese Licorice (Root) Glycyrrhiza glabra frish, Ashwagandha (Root) Withania somnifera, Rhodiola rosea (Root) 4:1 Extract

Amino | Fit Blend: 50 ma

Glutamine, Leucine, Isoleucine, Valine 2 mg

Delivery Agent: BioPerine®

† Daily Value (DV) not established

Other Ingredients: Gelatin, Magnesium Stearate, FD&C Blue 1, FD&C Red 40, FD&C Yellow 5, FD&C Yellow 6, Titanium Dioxide BioPerine® is registered trademark and patented product of Sabinsa

WARNING: KEEP OUT OF REACH OF CHILDREN. LipoRUSH® DS is intended for use by healthy individuals only. If you are taking any medications. planning any medical or surgical procedure or have any medical condition. consult your physician before use. Do not use LipoRUSH® DS or Censor® if

pregnant, nursing or have any medical condition. Use only as directed. Do not

DS contains caffeine. Do not use LipoRUSH® DS for more than 8 weeks. Consult with your physician prior to use if you are taking medication, including but not or epileptic disorders, difficulty urinating, diabetes, high or low blood pressure, cardiac arrhythmia, recurrent headaches, enlarged prostate or glaucoma. Discontinue 2 weeks prior to surgery or if you experience rapid heart beat, dizziness, severe headache, or shortness of breath. Do not use if you are prone to dehydration or exposed to excessive heat. Individuals who exceed the recommended serving or consume additional caffeine with this product may experience

## CENSOR

Supplement Facts

serving size 2 sortgets	bervings i er container 15	
	Amount Per Serving	% DV
Calories	20	
Calories from Fat	20	
Total Fat	2 g	3%**
Saturated Fat	0 g	0%**
Polyunsaturated Fat	1.5 g	†
Monosaturated Fat	0 g	†
Cholesterol	5 mg	2%
Protein	<1 mg	1%**
Vitamin E (as d-Alpha Tocopherol)	10 IU	33%
CLA/Omega 3-6-9 Bland:	2 000 mg (2 g)	+

Safflower Oil (Contains Conjugated Linoleic Acid (CLA), Tonalin® CLA) Omega-Krill Blend [Fish Oil (Contains Eicosapentaenoic Acid {EPA} Docosahexaenoic Acid (DHA)), Krill Oil), Flaxseed Oil (Linum usitatissimum) (Contains Alpha Linolenic Acid (ALA), Linoleic Acid (LA)), Evening Primrose Oil (Oenothera biennis)(Contains Linoleic Acid (LA), Gamma Linolenic Acid (GLA)), Borage Oil (Borago officinalis) (Contains Linoleic Acid {LA}, Gamma Linolenic Acid {GLA})

\*\*Percent Daily Values (DV) are based on a 2,000 calorie diet · Daily Value (DV) not established

Other Ingredients: Gelatin, Vegetable Glycerin, Mixed Natural Tocopherols. Contains Fish (Anchovy, Mackerel, Sardine, Shellfish (Krill)) Ingredients. Tonalin® CLA is an exclusively licensed trademark of Cognis under U.S. Patents #5,554,646, 5,428,072, 5,430,066 and 6,015,833.

## SUGGESTED USE FOR THE STACK:

Take 1 LipoRUSH® DS capsule daily, preferably with meals and 6-8 oz. of juice or water. To avoid sleeplessness, avoid taking LipoRUSH® DS within 6 hours of bedtime. Take 2 Censor® softgels up to three times daily with meals. To be used as part of a healthy lifestyle consisting of regular physical activity and a proper diet.

exceed recommended dose. Do not exceed recommended daily intake. Not intended for use by persons under age 18. Do not consume Synephrine or caffeine from other sources, including but not limited to, coffee, tea, soda and other dietary supplements or medications containing phenylephrine or caffeine. LipoRUSH® limited to MAOI inhibitors, anti-depressants, aspirin, nonsteroidal anti-inflammatory drugs or products containing phylephrine, ephedrine, pseudoephedrine, or other stimulants. Consult your physician prior to use if you have a medical condition, including but not limited to, heart, liver, kidney, or thyroid disease, psychiatric adverse side effects. Do not use if tamper resistant seals are broken. STORE IN A COOL DRY PLACE, AVOID EXCESSIVE HEAT.





✓ NOW WITH MORE ENERGY



Flaxseed Oil