

Human Lactic Commensals (HLC) probiotic Human Lactic Commensals (HLC) problotic strains were studied in a randomized, double-blind, placebo-controlled human trial at 10 billion CFU per day. Proprietary non-pathogenic human-sourced microflora bacterial strains are utilized as they have strong epithelial adherence and a naturally high tolerance to stomach acid.

Recommended intake: Children (1-4 years): In a glass, add water or milk to one scoop of HLC Baby F and mix. Take once daily with a meal, at least 2-3 hours before or after taking antibiotics, or as professionally directed.

Risk information: If you have an immune-compromised condition, do not use. If you are experiencing fever, vomiting, bloody damhea or severe abdominal pain, consult your healthcare practitioner prior to use. If symptoms of digestive upset occur, worsen or pensist beyond three days, discontinue use and consult your healthcare practitioner.

Vacuum-sealed for your protection and for product freshness. Do not use if outer seal is missing or broken. Store under refrigeration (below 46°F).

Keep out of reach of children

HLC BABY F

probiotic supplement

for formula fed babies





Pack size: Net Wt 2.3 oz (66 g)

Supplement Facts

Serving Size 1 Scoop (1.1 g) Servings Per Container 60

Amount per serving Total Carbohydrate HLC Consortium 10 hillion viable cells

Bifidobacterium bifidum (CUL-20) Bifidobacterium animalis subsp. lactis (CUL-34)

Lactobacillus salivarius (CUL-61) Lactobacillus paracasei (CUL-08)

† % Daily Value not established

Contains: Milk

Made in the HK for Seroval USA, Pittsburgh, PA 15275, 1-888-737-6925

Supplement Facts

HLC Consortium 10 hillion viable cells

Bifidobacterium bifidum (CUL-20) Bifidobacterium animalis subsp. lactis (CUL-34) Lactobacillus salivarius (CUL-61) Lactobacillus paracasei (CUL-08

Human Lactic

Commensals (HLC) probiotic strains were studied in a randomized, double-blind. placebo-controlled human trial at 10 billion CFU per day. Proprietary non-pathogenic human-sourced microflora bacterial strains are utilized as they have strong epithelial adherence and a naturally high tolerance to stomach acid.

HLC BABY F probiotic supplement

for formula fed babies





Recommended intake: Children

(1-4 years): In a glass, add water or milk to one scoop of HLC Baby F and mix. Take once daily with a meal, at least 2-3 hours before or after taking antibiotics, or as professionally directed.

Risk information: If you have an immune-compromised condition. do not use. If you are experiencing fever, vomiting, bloody diarrhea or severe abdominal pain, consult your healthcare practitioner prior to use. If symptoms of digestive upset occur, worsen or persist beyond three days, discontinue use and consult your healthcare practitioner. Safety-sealed for your protection and for product freshness. Store under refrigeration (below 46°F).

Keep out of reach of children



Gluten free

Made in the UK for Seroyal USA Pittsburgh, PA 15275, 1-888-737-6925

RARCODE LOCATION 883196 21660

Pack size: Net Wt 2.3 oz (66 g)