



LIPO CELL

Fat Burner*

Plus

HERBAL SUPPLEMENT

SLIM4life™

90 CAPSULES

Supplement Facts

Serving size 2 capsules

Servings per container 60

Amount per 2 capsules	% Daily Value
Proprietary blend 766mg	
Green Tea (leaf, extract 10:1)	†
Bitter Orange (immature fruit, extract 100:1)	†
Sweet-scented Geranium (flower oil, extract 50:1)	†

† Daily Value not established.

Other ingredients: Gelatin, cellulose, magnesium stearate, silicon dioxide, FD & C Red No. 3, FD & C Yellow No. 6, titanium dioxide.

Manufactured for: 4Life TM Holdings, LLC,
17470 North Pacesetter Way, Scottsdale, AZ 85255



***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.**

PLS1234120957V1 Lot No.:/Exp.:

Bitter Orange (*Citrus aurantium*) contains small amounts of alkaloids such as synephrine and octopamine, which are direct- and indirect-acting adrenergic agents. Sweet-scented Geranium oil (from *Pelargonium graveolens*) contains methylhexanamine, which is also a direct and indirect-acting adrenergic agent. Adrenergic agents with beta-agonist activity stimulate lipolysis and increase the resting metabolic rate in some persons, thus increasing the removal and oxidation of fat from adipose tissue stores. This action is potentiated by methylxanthines, such as caffeine, present in Green Tea (*Camellia sinensis*).*

DIRECTIONS: Take one (1) or two (2) capsules one (1) to three (3) times per day with an 8 oz (240 mL) glass of water, 30 minutes before meals. Commence with a low dose and build up to the most comfortable dosage level.

DO NOT EXCEED A DOSAGE OF 6 CAPSULES PER DAY OR TAKE MORE THAN TWO CAPSULES AT A TIME. WAIT AT LEAST 4 HOURS BETWEEN DOSES.

SALE TO, OR USE BY, PERSONS UNDER THE AGE OF 18 MUST HAVE PARENTAL APPROVAL. KEEP OUT OF THE REACH OF CHILDREN!

STORE AT ROOM TEMPERATURE (59°F - 86°F) IN A DRY PLACE AND KEEP TIGHTLY CLOSED.

Seek advice from a health care practitioner prior to use if you are pregnant or nursing, or if you have high blood pressure, heart or thyroid disease, diabetes, or if taking prescription drug. Do not take this product if you are taking appetite suppressants or products containing ephedrine or pseudoephedrine, or are taking an MAO inhibitor, or have taken an MAO inhibitor within the last 3 weeks. Reduce or discontinue use if nervousness, tremor, sleeplessness, severe loss of appetite or nausea occur.