201910-120

Cautions and Warnings: If you have gastrointestinal lesions/ulcres, bleeding disorted, responsible properties, or are taking anticoagulant / blood thinners, and i-inflammatory or antibiotic medications, rear about to have surgery, consult a healthcare rearbout to have surgery, consult a healthcare practitioner prior to use. Do not use if you are pregnant or breast-feeding. If you experience nausea, vomiting or hypersensitivity / allergic reaction while taking Wobenzym, discontinue use and consult vour healthcare practitioner.

Recommended dose: Adults: Take 2 tablets twice daily at least 45 minutes before meals with water.

Store in a cool dry place. Do not use if safety seal is broken or missing. Keep out of reach of children.

Distributed by: Mucos USA 112 Technology Drive Pittsburgh, PA 15108 Wobenzym.com Dietary Supplement

Wobenzym® plus

Helps to maintain healthy joints, mobility and flexibility¹

Temporary relief from everyday aches, pains and muscle soreness¹

Speedy recovery following exertion¹



120 enteric-coated tablets

¹ These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease

Supplement Facts

Serving Size 2 Tablets • Servings per container 60

		Daily alue
Bromelain** (from pineapple (Ananas comosus var comosus) stem)	270 mg (1350 FIP***-units)	
Trypsin (from porcine (Sus scrofa) or bovine (Bos taurus) pancreas)	144 mg (4320 FIP*** - units)	*
Rutoside trihydrate (Rutin) (from Styphnolobium japonicum	300 mg	1

- * Daily Value not established.
- ** From dried purified aqueous extract.
- *** FIP is the measurement of enyzme activity according to the methods of the International Pharmaceutical Federation.

Other ingredients: Microcrystalline cellulose, calcium phosphate, hydroxypropyl cellulose, vegetable stearate, enteric coating (sodium alginate, medium chain triglycerides, ethylcellulose, stearic acid, oleic acid, purified water) purified water.