

Warnings: Consult a healthcare professional if a sudden change in bowel habits occurs and persists for two weeks. Rectal bleeding or failure to have a bowel movement after use of a laxative may indicate a serious condition; discontinue use and consult a healthcare professional. Do not use if pregnant or nursing. Unless directed by a healthcare professional, do not use laxative products when abdominal pain, nausea, or vomiting is present, or if you have kidney disease. Do not take more than three tablets in a 24-hour period or use the maximum dosage for more than two weeks. If taking prescription drugs or if you have gallstones or bile duct obstruction, consult your healthcare professional prior to use.

For more information, please visit integrativepro.com.

Contains no sugar, salt, yeast, wheat, gluten, dairy products, artificial colors, flavors, preservatives, or ingredients of animal origin.

©2015 INTEGRATIVE THERAPEUTICS, LLC
GREEN BAY, WI 54311 USA
integrativepro.com • 800.931.1709

◆ LZ78967.A02 BLK896A



THIS STATEMENT HAS NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION.
THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.



LAXATIVE FORMULA

35 TABLETS

DIETARY SUPPLEMENT

Supplement Facts

Serving Size 3 tablets

Servings per container 11

Amount per 3 tablets	%DV
Calories	10
Total Carbohydrate	1 g <1%†
Dietary Fiber	1 g 4%†
Magnesium Hydroxide	2.6 g **
Proprietary Mucilage Blend: marshmallow (<i>Althaea officinalis</i>) root extract, slippery elm (<i>Ulmus rubra</i>) bark, and fennugreek (<i>Trigonella foenum-graecum</i>) seed extract	650 mg **
Burdock (<i>Arctium lappa</i>) Root Extract	300 mg **
Proprietary Soothing Blend: fennel (<i>Foeniculum vulgare</i>) seed, ginger (<i>Zingiber officinale</i>) rhizome extract, and peppermint (<i>Mentha x piperita</i>) leaf	150 mg **
Red Clover (<i>Trifolium pratense</i>) Blossom Extract	100 mg **

†Percent Daily Values (DV) are based on a 2,000 calorie diet.

**Daily Value not established.

Other ingredients: cellulose, stearic acid, modified cellulose gum, modified cellulose, magnesium stearate, glycerin, silicon dioxide, soy lecithin

Recommendation: Day 1 through day 7: Take 3 tablets at bedtime with 8 ounces of water. Day 8 through day 14: Take 2 tablets at bedtime with 8 ounces of water, or as recommended by your healthcare professional. Reduce dosage if you experience abdominal discomfort or diarrhea. If discomfort persists, discontinue use.