

Nutrition Facts

Serving Size: 1 Caplet

Servings per container: 100

Amount Per Serving: % Daily Value

Total Soluble Vitamins

Vitamin A (retinol + acetate)

2000 IU 40%

Vitamin E (alpha-tocopherol acetate)

30 IU 100%

Water-Soluble Vitamins

Vitamin B (ascorbic acid)

500 mg 833%

Vitamin B1 (thiamine mononitrate)

20 mg 1333%

Vitamin B2 (riboflavin)

20 mg 100%

Niacin (nicotinic acid)

500 mg 500%

Vitamin B6 (pyridoxine HCl)

25 mg 1750%

Biotin

100 mcg 30%

Pantothenic acid (D-calcium pantothenate)

25 mg 250%

Folic acid

1.0 mg 250%

Vitamin B12 (cyanocobalamin)

50 mcg 833%

Cobalamin (B12)

50 mcg

Minerals

Iron (ferrous fumarate)

27 mg 150%

Chromium (chromium chloride)

83 mcg 83%

Magnesium (magnesium oxide)

50 mg 12.5%

Manganese (manganese gluconate)

5 mg 250%

Copper (copper oxide)

3 mg 150%

Selenium (L-selenomethionine)

50 mcg 71%

Zinc (zinc oxide)

22.5 mg 150%

Daily Value (DV) not established

OTHER INGREDIENTS:

Microcrystalline Cellulose, Croscarmellose Gum, Fumed Silica,

Dicalcium Phosphate, Magnesium Stearate, Hydroxypropyl Methylcellulose, Polysorbate 80, FD&C Red #40 Aluminum Lake, Titanium Dioxide, FD&C Yellow #6 Aluminum Lake, FD&C Blue #2 Aluminum Lake, Titanium Dioxide.

DESCRIPTION

BACMIN is a prescription and multivitamin/mineral caplet specially formulated for prophylactic or therapeutic nutritional supplementation in physiologically stressful conditions.

BACMIN supplies therapeutic levels of water-soluble vitamins (ascorbic acid and B-vitamins), supplemental levels of fat-soluble vitamins (vitamins A and E), and minerals (iron, chromium, manganese, copper, selenium and zinc), plus magnesium.

INDICATIONS*

BACMIN is indicated for prophylactic or therapeutic nutritional supplementation in physiologically stressful conditions. These may include:

Conditions causing depletion, reduced absorption or bioavailability of essential vitamins and minerals

• Insufficient intake due to highly restricted or unbalanced diets such as those frequent associated with anorectic conditions and other states of severe malnutrition.

• Gastrointestinal disorders, chronic alcoholism, chronic or acute infections (especially if involving the ileum), prolonged or wasting disease, congestive heart failure, hyperthyroidism, poorly controlled diabetes or other physiologic stress.

• Also, patients on estrogen and corticosteroid or other estrogen therapy, anti-hepatitis which affect intestinal muciniferous, or other interfering drugs.

Certain conditions resulting from severe B-vitamin or ascorbic acid deficiency

• Recovery from surgery or trauma involving severe burns, fractures or other extensive damage.

• Also, pregnant women and those with heavy menstrual bleeding.

*This statement has not been evaluated by the Food and Drug Administration.

This product is not intended to diagnose, treat, cure, or prevent disease.

CONTRAINDICATIONS

BACMIN is contraindicated in patients hypersensitive to any of its components.

WARNINGS

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Not intended for treatment of pernicious anemia or other megaloblastic anemias where vitamin B12 is deficient. Neurological involvement may develop or progress. Despite temporary relief of anemia, patients with vitamin B12 deficiency who receive supplemental folic acid are unduly exposed to treatment with B12.

PRECAUTIONS

General: DO NOT EXCEED RECOMMENDED DOSE.

Folic Acid: Folic acid in doses above 5 mg daily may obscure pernicious anemia in that hematology reactions can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask symptoms of pernicious anemia.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Drug and Treatment Interactions

Adverse reactions have been reported with specific vitamins and minerals, but generally at substantially higher doses than in BACMIN. However, drug-drug interactions may occur at lower levels. Iron, even at the usual recommended level, has been associated with gastrointestinal constipation in some patients.

ADVERSE REACTIONS

Adverse reactions have been reported with specific vitamins and minerals, but generally at substantially higher doses than in BACMIN. However, drug-drug interactions may occur at lower levels. Iron, even at the usual recommended level, has been associated with gastrointestinal constipation in some patients.

MARNEL

0682-3001-01

BACMIN

CAPLETS

VITAMIN/MINERAL NUTRITIONAL
SUPPLEMENT

Rx

100 CAPLETS

Manufactured for:
MARNEL PHARMACEUTICALS, LLC
Lafayette, Louisiana 70506

KEEP OUT OF REACH OF CHILDREN

Tamper Evident: Do not use if seal is broken or missing.
See attached labeling for complete product information.

MADE IN CANADA

Rev. 01/75



WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

DIRECTIONS: Adults: Take one caplet daily, or as prescribed by a licensed medical practitioner.

Children under 12: Consult a doctor.

STORAGE: Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F). [See USP Controlled Room Temperature.] Protect from light and moisture.