GRG

Potencies verified by GNC procedures #5103 and #5118. Conforms to USP <2091 > for weight. Meets USP <2040> disintegration.

\* This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

KEEP OUT OF REACH OF CHILDREN. Store in a cool, dry place,

For More Information: 1-888-462-2548

## SHOP NOW @ GNC.COM

Distributed by: **General Nutrition Corporation** Pittsburgh, PA 15222 USA



## GNC

## **B-Complex**

**50 MG** 

Essential for energy production & carbohydrate metabolism\*

**DIETARY SUPPLEMENT** 100 CAPSULES

CODE 017914

Directions: As a dietary supplement, take one capsule daily.

## **Supplement Facts**

Serving Size One Capsule

| Amount Per Serving                         | % Dai      | ly Value |
|--|------------|----------|
| Thiamin (Vitamin B-1)(as Thiamin Mononitra | ate) 50 mg | 3333%    |
| Riboflavin (Vitamin B-2)                   | 50 mg      | 2941%    |
| Niacin (as Niacinamide)                    | 50 mg      | 250%     |
| Vitamin B-6 (as Pyridoxine Hydrochloride)  | 50 mg      | 2500%    |
| Folic Acid                                 | 400 mcg    | 100%     |
| Vitamin B-12 (as Cyanocobalamin)           | 50 mcg     | 833%     |
| Biotin                                     | 50 mcg     | 17%      |
| Pantothenic Acid (as Calcium d-Pantotheni  | ate) 50 mg | 500%     |
| Choline Bitartrate                         | 50 mg      | *        |
| Inositol                                   | 50 mg      | *        |
| para-Aminobenzoic Acid (PABA)              | 50 mg      | *        |

\* Daily Value not established.

Other Ingredients: Gelatin, Mono- & Diglycerides, Whole Brown Rice Powder (Orvza sativa), Brewer's Yeast, Soy Lecithin, Alfalfa Leaf Powder (Medicago sativa). Wotercress Herb Powder (Nasturtium officinale), Porsley Herb Powder (Petroselinum crispum). CONTAINS: Wheat

WARNING: Consult your physician prior to using this product if you are pregnant, nursing, taking medication, or have a medical condition. Discontinue use two weeks prior to surgery.