

Each serving provides 200 mg of caffeine.

KEEP OUT OF REACH OF CHILDREN.

Store in a cool, dry place.

For More Information:

1-888-462-2548

SHOP NOW @ GNC.COM

Distributed by:

General Nutrition Corporation

Pittsburgh, PA 15222

^See carton for related information.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Capsimax® trademark belongs to OmniActive Health Technologies.



GNC

TOTAL LEAN™
ADVANCED

METABOLIC ELITE™

DIETARY SUPPLEMENT

FAST-ACTING METABOLISM
SUPERCHARGER

HIGH
ENERGY
THERMOGENIC

CLINICALLY PROVEN TO
BURN 6X
MORE CALORIES!™

STIMULATES
**MIND
& BODY**

CLINICALLY STUDIED
60 TABLETS

CODE 489212

CRG

DIRECTIONS: As a dietary supplement, take two fast-acting tablets 30 minutes prior to exercise. Do not take near bedtime.

Supplement Facts

Serving Size Two Fast-Acting Tablets

Servings Per Container 30

Amount Per Serving	% Daily Value
Niacin	20 mg 100%
6x Clinical Burn Blend	
Mucuna pruriens Seed Extract	500 mg *
(20% L-Dopa = 100 mg)	
Ginger Root Extract (Zingiber officinale)	500 mg *
(5% Gingerol = 25 mg)	
Boswellia serrata Gum Extract	255 mg *
(60% Boswellic Acid = 153 mg)	
Caffeine Anhydrous	200 mg *
Capsimax® Capsicum Seed Extract (Capsicum annuum)	100 mg *
Cinnamon Bark Extract	100 mg *
(Cinnamomum zeylanicum)(15% Polyphenols = 15 mg)	
Piper longum Fruit Extract	2 mg *
Botanical Rush Blend	
Green Coffee Bean Extract	200 mg *
(Coffea canephora)(45% Chlorogenic Acid = 90 mg)	
Rocket Plant Leaf Extract (Eruca sativa) &	100 mg *
Chicory Root Extract (Cichorium intybus) Blend	
Mind/Body Matrix	
Reishi Mushroom Powder (Ganoderma lucidum)	90 mg *
Cordyceps sinensis Extract	10 mg *

* Daily Value not established.

OTHER INGREDIENTS: Cellulose, Titanium Dioxide (Natural Mineral Whitener), Vegetable Acetoglycerides, FD&C Blue #1 Aluminum Lake, Riboflavin (Color).

WARNING: Consult your physician prior to using this product if you are pregnant, nursing, taking medication, or have a medical condition. Discontinue use two weeks prior to surgery.