

[Kroger Ginkgo biloba capsules: \(Size: 120 × 100 mg capsules / bottle\)](#)



Ginkgo Biloba Soft Capsule Product Specifications: 120 × 100 mg capsules / bottle

[Kroger Ginkgo biloba capsules: \(Size: 120 × 100 mg capsules / bottle\) - American products](#)

[Kroger Ginkgo biloba capsules: \(Size: 120 × 100 mg capsules / bottle\)](#)

[Name] Kroger Ginkgo biloba capsules

[English name] Kroger Ginkgo Biloba

The main components each containing (24% flavonol glycosides) 120 mg

[Specification] 120 × 100 mg capsules / bottle

[Suggested Use] Adults: one day, with a meal

Memory disorders and cardiovascular function: Under normal circumstances, 120 mg.

If more severe dementia or Alzheimer's disease at doses up to 240 mg per day may be necessary.

Intermittent claudication: 120 per day - 240 mg.

Glaucoma: 120 mg per day have improved vision.

[Note] Preliminary results often take 4 - 6 weeks, beyond this period, the effect should be enhanced.

Supplement Facts

Serving Size: 1 Capsule
Servings Per Container: 100

	Amount Per Serving	% Daily Value (DV)
Ginkgo Biloba Extract (Ginkgo biloba) (leaf) (standardized to contain 24% ginkgolide flavonols) (9500000)	120 mg	**
Ginkgo biloba (Ginkgo biloba) (leaf)	100 mg	**

Other Ingredients: Rice Powder, Gelatin Capsules, Vegetable Magnesium Stearate, Silica.

[] Other ingredients [rice flour](#), [gelatin capsules](#), [vegetable magnesium stearate](#), [silica](#). Ginkgo (Ginkgo biloba) is one of the oldest species, its leaves are today the most widely used herbs. In Europe and the United States, [ginkgo biloba supplements](#) are the best-selling herbs. It is always ranked in the regulations of France and Germany's top medicine. Click on the following link directory [ginkgo biloba supplements](#) doctor online to provide you with more detailed information:

- [Ginkgo biloba supplements](#)

WARNING [epilepsy](#), people should not take [ginkgo](#), because the fear that it may lead to seizures.

[Pregnant and lactating women](#) should not take [ginkgo](#).

People who have diabetes before taking [ginkgo](#) should ask their doctor.

Because of the risk of bleeding before surgery or dental surgery at least 36 hours you should stop taking it.

[Storage] store in a cool, place dried reach of children.

[Validity] is usually 24 months, see a bottle of validity date