Supplement Facts Serving Size: 1 bottle (10 fl. oz.) Servings Per Case: 12

Amount Per Serving		% DV
Calories	8	
Total Carbohydrate	2g	<1%**
Sugars	0g	1
Niacin (as Niacinamide)	65mg	325%
Vitamin B6 (as Pyridoxal-5-Phosphate)	650mcg	33%
Folic Acid	325mcg	81%
Vitamin B12 (as Methylcobalamin)	45.5mcg	758%

Explosive Energy Blend CarnoSyn® Beta-Alanine, Arginine Nitrate (NO3-T®), Caffeine Anhydrous (200mo), N-Acetyl-L-Tyrosine

**Percent Daily Values (% DV) are based on a 2.000 calorie die † Daily Value not established

Other Ingredients: Water, Tartaric Acid, Natural and Artificial

Flavors, Sucralose, Acesulfame Potassium, FD&C Red #40

Made in the U.S.A. using U.S. and Imported Ingredients and Components

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure

NO3-T® is a trademark of ThermoLife International, LLC.

CarnoSym Datents as listed on www.carnosyn.com and registered

Cellucor®, C4® and G4 Chrome Series™ are trademarks

nutrabolt*

Nutrabolt® 3891 S. Traditions Dr., Bryan, TX 77807 1.866.927.9686 - www.cellucor.com



SUGGESTED USE:

Consume one bottle (10 fl. oz.) of C4® ON THE GO 20-30 minutes before dominating exercise . . and life. Some individuals may experience a harmless tingling sensation, which is attributed to beta alanine.

HEALTHY ADULTS 18 YEARS OF AGE OR OLDER Do not us this product if you are pregnant, nursing, or are currently taking nitrates for chest pain or if you are taking medication used to treat erectile dysfunction such as PDF-5 inhibitors Before using this product, consult a licensed, qualified health care professional, including but not limited to, if: you are taking antidepressants such as a MAOI (Mongamine anti-inflammatory drugs, pseudoephedrine, or you are taking over-the-counter medication; or if, you suspect you have o have been treated for, diagnosed with or have a family history any medical condition, including but not limited to: high o low blood pressure, diabetes, glaucoma, anxiety cardiovascular, psychiatric or seizure disorders, cardiac arrhythmia, stroke, heart, liver, kidney or thyroid disease, o and medications. Discontinue use 2 weeks prior to surgery Immediately discontinue use and contact a medical doctor in you experience any adverse reaction to this product. Do no use if safety seal is broken or missing. Store in a cool dry place. Do not Freeze. KEEP OUT OF REACH OF CHILDREN.

