







Cellucor® and C4® are trademarks of and distributed by:



Nutrabolt® 3891 S. Traditions Dr. Brvan, TX 77807 1.866.927.9686 - www.cellucor.com 103408

CELLUCOR

Supplement Facts Serving Size: 1 bottle (10 fl. oz.) Servings Per Container: 4

Amount Per	Serving	% DV
Calories	8	
Total Carbohydrate	2g<	< 1%**
Sugars	0g	
Vitamin B6 (as Pyridoxal-5-Phosphate)45 mcg	2%
Vitamin B12 (as Methylcobalamin)	45 mcg	750%
Phosphorus (as DiSodium Phosphate & DiPotassium Phosphate)	133 mg	13%
Magnesium (as Magnesium Chloride)	30 mg	8%
Chloride (as Magnesium Chloride)	99 mg	3%
Sodium (as DiSodium Phosphate)	160 mg	7%
Potassium (as DiPotassium Phosphate	e) 60 ma	2%

C4® Sport Blend

4179ma Beta Alanine, Arginine Alpha-Ketoglutarate, Taurine, Caffeine Anhydrous (180 mg)

**Percent Daily Values (% DV) are based on a 2,000 calorie die † Daily Value not established

Other Ingredients: Water, Malic Acid, Artificial Flavors, Sucralose, Acesulfame Potassium, FD&C Blue #1

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

SUGGESTED USE: Consume one bottle (10 fl. oz.) of C4® SPORT ON THE GO 20-30 minutes before dominating exercise...and life. Do not consume more than 2 bottles of C4® SPORT ON THE GO per day. Some individuals may experience a harmless tingling sensation, which is attributed to beta alanine. WARNING: THIS PRODUCT IS ONLY INTENDED TO BE CONSUMED BY HEALTHY ADULTS, 18 YEARS OF AGE OR OLDER. Do not use this product if you are pregnant or nursing. Before using this product, consult a licensed, qualified, health care professional, including but not limited to, if: you are taking antidepressants such as a MAOI (Monoamine Oxidase Inhibitor) or SSRI, blood thinners, nonsteroidal anti-inflammatory drugs, pseudoephedrine, or you are taking any other dietary supplement, prescription drug or over-the-counter medication; or if, you suspect you have or have been treated for, diagnosed with or have a family history of, any medical condition, including but not limited to: high or low blood pressure, diabetes, glaucoma, anxiety, cardiovascular, psychiatric or seizure disorders, cardiac arrhythmia, stroke, heart, liver, kidney or thyroid disease, or difficulty urinating due to prostate enlargement. This product contains caffeine and should not be used by individuals wishing to eliminate caffeine from their diet or in combination with caffeine or stimulants from other sources, including but not limited to, coffee, tea, soda, or other dietary supplements and medications. Discontinue use 2 weeks prior to surgery. Immediately discontinue use and contact a medical doctor if you experience any adverse reaction to this product. Do not use if safety seal is broken or missing. Store in a cool dry place. KEEP OUT OF REACH OF CHILDREN. ‡ Based on sales of C4® brand products in the U.S.A. from 2014 to the present.